Eye of the Eagle

Volume 12, Number 1 | THE CARTER CENTER | February 2011

The number of people at risk for onchocerciasis (river blindness) in the Americas has declined for the first time since The Carter Center began its campaign to eliminate the disease in the Western Hemisphere, according to data presented at the 20th annual Inter-American Conference on Onchocerciasis (IACO) in Antigua, Guatemala, Nov. 10–12, 2010. In 1993, the estimated at-risk population was 4 million. It is now fewer than 500,000.

Most recently, the at-risk population has declined by nearly

The current at-risk population includes only those living in the remaining 10 foci where the disease is either actively transmitted or where transmission has

What's Inside

ments. More than 9.2 million people were treated with Pfizer-donated Zithromax® for trachoma (see Figure 4), and 120,000 people were tested for malaria. Of those tested for malaria, 70,000 required and received treatment. Health workers noted that now that the Ethiopians are experienced and comfortable with the campaign, they are demanding service from the field teams, and supervision is coming as much from the program recipients as from the implementers.


1
MEXICO
3
GUATEMALA
5
6

Figure 2)RXU E7LPHV ESHUE<HDU 7UHDWPHQW ZLWK ,YHUPHFWLQ LQ WKH $PHULFDV 5HJLRQ
Projected for 2011



The 2010 IACO meeting was convened by the Ministry of Health of Guatemala, the Carter Center's Onchocerciasis Elimination Program for the Americas (OEPA), and the Pan American Health Organization, with support from the Bill & Melinda Gates Foundation, the Lions Clubs International Foundation, and Merck. It was attended by 75 people, including 25 representatives of the Guatemala Ministry of Public Health and Social Assistance, from national, departmental, and municipal levels. Dr. Salomón López, director of the Department of Health Regulation, Surveillance, and Control, represented the minister of health of Guatemala during the opening ceremony. Also in attendance were the directors of the six national onchocerciasis elimination programs in Brazil, Colombia, Ecuador, Guatemala, Mexico, and Venezuela; representatives of the local Lions Clubs from five of the six countries (Alfonso Barahona, Margarita Peña Constante, Ramiro Peña Constante, Dr. Libardo Bastidas Passos, Dr. Ricardo Araujo Gurgel, Vania Araujo Gurgel, and Dr. Florencio Cabrera Coello); Kristen Eckert (Lions Clubs International Foundation); Dr. Steven Ault (PAHO); Dr. Adrian Hopkins (Mectizan Donation Program); Ken Gustavsen (Merck); Dr. Ed Cupp (chair of OEPA's Program Coordinating Committee); and Dr. Mark Eberhard (U.S. Centers for Disease Control and Prevention). Attending for The Carter Center were Dr. Mauricio Sauerbrey (director of OEPA), Dr. Donald Hopkins (vice president for health programs), Dr. Frank Richards, Craig Withers, Lindsay Rakers, and Nicole Kruse.

ment approach is being used to hasten disease elimination in four foci: South Chiapas (Mexico), Northeast and South (Venezuela), and Amazonas (Brazil) (see Figure 2). Four-times-per-year treatment occurred in 2010 in all hyperendemic areas of South Chiapas and the Venezuelan Northeast and in 40 percent of Venezuelan South. In 2011, this treatment intensification will expand to 100 percent of hyperendemic areas in Brazil and the South focus of Venezuela. Several countries are also considering using prolonged (5–6 week) courses of doxycycline treatment for onchocerciasis as well.

The progress of the OEPA initiative was palpable at this meeting and is possible only due to the persistence, expertise, and creativity of the partners. One memorable presentation was given by an acting troupe that has provided "art as a bridge to health" services for health education through onchocereiasis fairs held in endemic communities. Troupe members provided lively and unforgettable performances of their unique methods of encouraging people to keep taking their Mectizan until onchocerciasis has been eliminated.

Caja Lúdica

A Guatemalan acting troupe demonstrates health education through song.

The Wadelai focus in Uganda became the first area in the country where a committee has recommended that mass drug administration with Mectizan® be halted, which means that experts

A

Figure 4 3HUVRQV 7UHDWHG IRU 7UDFKRPD $]LWKURP\
7HWUDF\FOLQH (\H 2LQWPHQW 'XULQJ 0DOWL

Five years ago in Ethiopia, 3-year-old girl Haymanot Shibabow charmed former U.S. President Jimmy Carter and former First Lady Rosalynn Carter with her willingness to demonstrate her training latrine. The Carters were visiting the girl's village, Mosebo, to help launch the expansion of trachoma control programming to the whole of Amhara region in September 2005.

Haymanot and her family were early adopters of improved sanitation practices in the region, and five years later they are still following good domestic hygiene. While her father has rebuilt the household latrine twice, young Haymanot still prefers to have her own, covered now with an old plastic basin.

The SAFE strategy (surgery, antibiotics, facial cleanliness, environmental improvement) has successfully been expanded to 150 districts in Amhara since 2005, and Mosebo has not

Health Workers Give Donated Zithromax, a First for Nigeria

With a quick swallow of banana-flavored medicine, David Nuhu, a 4-year-old boy from Aloshi village in central Nigeria, became the first recipient of Pfizer-donated Zithromax® for trachoma in the country.

Elizabeth Cromwell

With assistance from community drug distributor Zinariya Samson, 4-year-old David Nuhu, from Aloshi village in Nasarawa state, takes the first dose of donated Zithromax in Nigeria to treat trachoma.

Community drug distributors Zinariya Samson and Daniel Adeka treated Nuhu's entire household during training exercises, providing not only antibiotics for trachoma control but also health education to help prevent future trachoma infections.

"I want my community to be healthy," Samson said, adding that her sister previously had distributed medication in her community.

The new program was officially launched in a ceremony in Abuja on Oct. 14, 2010, and included representatives of the federal Ministry of Health, Pfizer Inc, and The Carter Center. The ceremony coincided with World Sight Day to raise awareness of preventable blindness in Nigeria. In addition to being Africa's most populous country, Nigeria is thought to account for a large share of the worldwide burden of trachoma. Nine northern states are endemic for the disease.

In 2010, Nigeria began implementing mass drug administration for trachoma control with Zithromax and tetracycline eye ointment in seven local government areas in Plateau and Nasarawa states. At the end of 2010, Sightsavers International also supported mass drug administration for trachoma in several areas in Zamfara, Sokoto, and Kebbi states.

In Plateau and Nasarawa, the Nigeria Ministry of Health and The Carter Center implement integrated drug administration for schistosomiasis, lymphatic filariasis, and river blindness control. Community drug distributors—like the two who treated David Nuhu's family—play a crucial role in the success of the program. Their participation maximizes the program's

continues on page 10

Global Health News

During the first 11 months of 2015, 25 countries reported 1,686 (95 percent) of 1,785 cases of Guinea worm disease reported in the world, while the other three endemic countries (Ethiopia, Ghana, and Mali) combined reported only 85 cases of the disease (se(uring)-8(nknow

Figure 5 'LVWULEXWLRQ F\ &RXQWU\ RI RQGLJHQRX
Cases of Dracunculiasis: 2010

coverage and reduces the cost of treatment. Communities are respon sible for providing an incentive to their community drug distributors.

Zinariya Samson will try to juggle her daily household chores with her duties as a community drug distributor during this treatment round. “Nothing good is easy,” she said.

In September, representatives from the Plateau and Nasarawa state ministries of health were trained in trachoma treatment protocol, report ing, supervision, and stock manage ment at the Carter Center office in Jos. The state ministry representatives then helped train integrated health teams from the seven local government areas where mass drug administration for trachoma is warranted. Participants included a representative from the Blindness Prevention Program in the federal Ministry of Health; represen tatives from Zamfara, Sokoto, and Kebbi states; staff from Sightsavers International Nigeria; and Christoffel-Blindenmission.

These representatives then returned home to lead training of community drug distributors in their local areas. From late September through October, 1,710 distributors were trained in Plateau and Nasarawa states; they will target some 780,000 people in the two states for treatment with antibiotics.

Community drug distributor Jacob Igbeadio, who has been doing the work for four years, said, “I feel an obligation to serve my village. Whenever I am distributing, they joke and call me ‘doctor.’ I like that.”



Patricia Graves