DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service

Centers for Disease Control

And Prevention (CDC)

Memorandum

Date: March 31, 2022

From: WHO Collaborating Center for Dracunculiasis Eradication, CDC

Subject: GUINEA WORM WRAP-UP #286

To: Addressees

"*Just 15* (video):

https://video.cartercenter.org/Default.aspx?youtube_id=PQZjCteVvFA&category=Health&filter=Guinea%20Worm%C2%A0Eradication%20Program

Figure 1



Ms. Faustina Fynn-Nyame of the C

INTERNATIONAL GW REVIEW MEETS VIRTUALLY; 15 CASES IN 2021

About 110 persons participated in the 25th International Review Meeting of Guinea Worm Eradication Program Managers organized by The Carter Center in cooperation with the World Health Organization that met virtually on March 9-11, 2022. Carter Center Chief Executive Officer Paige Alexander and World Health Organization Director-General Dr. Tedros Ghebreyesus greeted participants in recorded videos. National Program Coordinators from the five endemic countries (Angola, Chad, Ethiopia, Mali, South Sudan), two pre-certification countries (Democratic Republic of the Congo, Sudan) and one country with cross-border infections

The new Minister of Public Health and National Solidarity Dr. Abdoulmadjid Abderahim and the Secretary-General Dr. Ishmael Barh Bachar both addressed the review meeting, in addition to National Program Coordinator Dr. Ouakou TCHINDEBET, who presented Chad's report. With

South Sudan GWEP Surveillance Snapshot 2021
Accessibility: 100%
Villages reporting 1+ GW infection: 4
Number of districts by surveillance level: 4 in level 1; 10 in level 2; 66 in level 3
Villages under Active Surveillance (VAS): 2,012 (1,401 level 1; 611 level 2)
Monthly reporting rate for VAS: 98%
Number of rumors: humans 48,041 (99% investigated in 24h), animals 548 (99% investigated in 24h)
Cash reward awareness: 84% humans, 68% animals
Cash reward amount: US$75 equivalent for reporting a confirmed human case, US$75 for reporting a confirmed infected animal
Integrated surveys: 1,045,357 persons and 27,894 animals screened for GW in case sweeps
Number and reporting rate for Integrated Disease Surveillance and Reporting (IDSR): 92%
% presumed sources of human cases identified*: 0% (0/4)
% human and animal Guinea worm infections contained: 25% (1/4)
*see definition page 11

MALI reported 2 human cases (50% contained) and 17 animal infections (16 dogs, 1 cat; 10/17 or 59% contained) in 2021 (see line list in issue #285). Mali's GWEP continues to educate dog vendors in Segou and Mopti Regions despite challenges of insecurity. It applies Abate to known or suspected contaminated water sources within seven days. The program began pilot-testing proactive tethering of dogs in two endemic villages of Djenne and Macina districts of Mopti and

ETHIOPIA

CAMEROON, which reported its last indigenous GW case in 1997 and was certified free of the disease by WHO in 2007, reported 2 human cases (0% contained), 15 infected dogs (67% contained) and 1 infected cat (uncontained) in Guere district which borders endemic Bongor district in Chad, in 2019-2021. The Guinea worm infections and cases in Cameroon are in a population of fishermen and farmers who share ethnicity, family links, and travel on both sides of the border (see discussion in *Guinea Worm WrapWorm Wrap*

Figure 3


Guinea W
100
Safe Water
79
Abate
92
Abate®
100
Containment
63
81
Safe Water
Abate®
10
Abate®
100
Con
tainm
Bury Fish Guts
NA
50
0
25
50
75
100
0
25
50
75
100
*See criteria for each i
text.
** The MGWEP encourages households to dry fish guts
them to chicken farmers.
NA: Not Applicable

RESEARCH REVIEW MEETING

Carter Center Associate Director for Research Dr. Fernando Torres-Velez, DVM, PhD, convened a virtual Research Review Meeting with over 70 participants on March 16, 2022, of researchers supported by The Carter Center on aspects of Guinea worm infections. Topics presented and discussed at the meeting included baboon surveillance in Ethiopia, miRNA biomarkers, simulation modeling of transmission dynamics, studies of Flubendazole for treating infections in dogs in Chad, Guinea worm genomics, copepod ecology, and remote sensing of surface water bodies.

MODIFIED INTERVENTION INDICES TO REFLECT VARIABLE MODES OF TRANSMISSION

With *D. medinensis* infections occurring in animals in three of the final four endemic countries (South Sudan is the exception) and evidence mounting to suggest that the infection is being transmitted to humans and animals not just by drinking water, as before, but likely also by people and animals eating raw or undercooked transport hosts such as small fish (up to 2-3.inches/5-7.5 cm long) and

The latter indicator, as first reported on in GW Wrap-Up #257, is intended to estimate GW programs' safe access to areas of greatest concern now for supervision and interventions. After transmission is interrupted nationwide, the entire country will need to be accessible for adequate surveillance and certification. Our first concern now, however, is to stop transmission, which requires safe access. The four main considerations for the new indicator are: 1) the denominator = surveillance level 1 (known or suspected endemic) plus option to include other areas if judged appropriate; 2) scores are 0 = not accessible for supervision and interventions, 1 = partly accessible, 2 = fully accessible; 3) administrative level= district or county; 4) all GW infections count, whether human or animal. Total score is sum of scores for all districts/counties of concern divided by maximal score (2x total number of districts/counties of concern) times 100 = percentage. A country's score may change with changes in security situations on the ground.

DEFINITION OF A PRESUMED SOURCE OF GUINEA WORM INFECTION

A presumed source/location of a human dracunculiasis case is considered identified if:

The patient drank unsafe water from the same source/location (specify) as another human case(s) or an infected domestic animal 10-14 months before infection, or

The patient lived in or visited the (specify) household, farm, village, or non-village area of (specify) a Guinea worm patient or an infected domestic/peri-domestic animal 10-14 months before infection, or

The patient drank unsafe water from (specify) a known contaminated pond, lake, lagoon or cut stream 10-14 months before infection.

If none of the above is true, the presumed source/location of the infection is unknown. Whether the patient's or animal's residence is the same as the presumed source/locality of infection or not should also be stated in order to distinguish indigenous transmission from an imported case.

DEFINITION OF A CONTAINED CASE*

A case of Guinea worm disease is contained if all of the following conditions are met:

1. The patient is detected before or within 24 hours of worm emergence; and
2. The patient has not entered any water source since the worm emerged; and
3. A village volunteer or other health care provider has properly managed the case, by cleaning and bandaging until the worm is fully removed and by giving health education to discourage the patient from contaminating any water source (if two or more emerging worms are present, the case is not contained until the last worm is pulled out); and
4. The containment process, including verification that it is a case of Guinea worm disease, is validated by a supervisor within 7 days of the emergence of the worm and
5. ABATE is used if there is any uncertainty about contamination of sources of drinking water, or if a source of drinking water is known to have been contaminated.

*The criteria for defining a contained case of Guinea worm disease in a human should be applied also, as appropriate, to define containment for an animal with Guinea worm infection.

RECENT PUBLICATIONS

Grunert R, Box E, Garrett K, Yabsley M, Cleveland C, 2022. Effects of temephos (Abate), Spinosad (Natular), and diflubenzuron on the survival of cyclopoid copepods. Am J Trop Med Hyg 106:818-822.

Note to contributors: Submit your contributions via email to Dr. Sharon Roy (gwwrapup@cdc.gov) or to Adam Weiss (adam.weiss@cartercenter.org), by the end of the month for publication in the following month's issue. Contributors to this issue were: the national Guinea Worm Eradication Programs, Dr. Donald Hopkins and Adam Weissof The Carter Center, Dr. Sharon Roy of CDC, and Dr. Dieudonné Sankara of WHO.

WHO Collaborating Center for Dracunculiasis Eradication, Center for Global Health, Centers for Disease Control and Prevention, Mailstop A-06, 1600 88 1 165y 0 4C0052 500912 0 612 792 reW 000912 0 612 h.q454168.26 474-5(n) 8wb63Gc*